

Medicinal Plant: (In Rajasthan)

| Botanical name | Family | Local Name | Part(s) Used | Ailments | Ethno Medicinal Uses |
|---------------------------------------|---------------|---------------|--------------|------------------------------------|--|
| <i>Abrus precatorius</i> Linn. | Leguminosae | Chirmi, Gunja | Whole plant | Wounds, polyurea, arthritis, fever | Used in healing, ear, abortifacient, polyurea and antifertility. |
| <i>Abutilon indicum</i> Linn. | Malvaceae | Jhumka | Leaf, seeds | Syphilis | The leaves and seeds are used to cure syphilis. |
| <i>Acacia catechu</i> (L.f.) Willd. | Leguminosae | Khair | Whole plant | Dental pain, | Abortifacient, tooth ache and gonorrhoea. |
| <i>Acacia ferruginea</i> D.C. | Leguminosae | Ker | leaf, fruit | Hypertension | Used for ear disorders and otorrhoea |
| <i>Acacia leucophloea</i> Willd | Leguminosae | Safed kikar | Leaves, bark | Urinary tract bleeding | To increase sexual potency, as a contraceptive and for menstrual complaints. |
| <i>Acacia nilotica</i> Willd. | Leguminosae | Babul | Whole plant | Sexual, burning | Used for burning sensations, in the eyes and asthma. |
| <i>Achyranthes aspera</i> Linn. | Amaranthaceae | Aandhi jhara | Whole plant, | Pneumonia, headache, earache | For headache, and ear-ache |
| <i>Actiniopteris dichotoma</i> Bedd. | Polypodiaceae | Morpankhi | Whole plant | Fever | Use to control typhoid and fever. |
| <i>Adiantum capillusveneris</i> Linn. | Polypodiaceae | Hansraj | Leaf | Leucorrhoea | Paste of leaves is used in sweta pradar. |
| <i>Aegle marmelos</i> (Linn.) Corr. | Rutaceae | Bel | Fruit | Pain, diarrhoea, | For sexual debility, diarrhoea and |

| | | | | | |
|--|-----------------|-----------|-------------|-------------------------|---|
| | | | | | constipation. |
| Alangium salvifolium (L. f.) Wang. | Alangiaceae | Aankol | Root | Abortificiet | Root extract and juiceisan abortifaicient |
| Allium cepa Linn. | Liliaceae | Pyaj | Bulb | Cholera | Used for cholera. |
| Allium Sativum Linn. | Liliaceae | Lahasun | Bulb | Ringworm | Applied to the affected part. |
| Aloe barbadensis Mill. | Liliaceae | Gawarpata | Leaves | Indigestion, rheumatism | Fresh pulp is used for indigestion, burns, Rheumatism, liver diseases and wounds. |
| Amaranthus spinosus Linn. | Amaranthaceae | Chandalai | Leaf | Urinary disorder. | Leaves are used orally as a diuretic. |
| Amorphophallus paconifolius (Dennst.) | Araceae | Jamikand | Rhizome | Hyper acidity | Paste of rhizomes is used orally with water. |
| Ampelocissus latifolia (Roxb.) Planch. | Vitaceae | Tita | Tuber | Fractured, Tuberculosis | To cure bone fractures, dyspepsia, indigestion and tuberculosis orally. |
| Anethum graveolens Linn. | Umbelliferae | Swa | Seed | Lactogogues | To digest food and secretion of milk. |
| Anogeissus latifolia Bedd. | Combretaceae | Dhokada | Gum | Back pain | Used to cure the damaged tissue during delivery. |
| Argemone Mexicana Linn. | Papaveracea | Dhaturi | Stem, latex | Rheumatism, Eye | Applied to affected parts, poultice for the eyes. |
| Arisaema tortuosum (Wall.) Schott | Aracaceae | Halida, | Tuber | Snake bite | Snake bite, bone fracture in animals. |
| Aristolochia bracteata Retz. | Aristolchiaceae | Kalipad | Whole plant | Boils, Stiff hair | Applied for boils and snake bite. |

| | | | | | |
|---------------------------------|---------------|----------|--------------|--------------------------------|---|
| Asparagus racemosus Wild. | Liliaceae | Satawari | Root | Debility, Impotency | Increased lactation, diabetes, muscle pain, muscular disorders, leucorrhoea, fertility. |
| Atylosia scarabaeoides Benth. | Leguminosae | Valoli | Root | Diarehoea | Juice of the root is used orally for diarrhea. |
| Azadirachta indica A. Juss. | Miliaceae | Neem | Seeds, leaf | Malaria, Worm | Used for excessive bleeding, as a contraceptive |
| Balanites aegyptiaca Linn. | Simarubaceae | Hingora | Whole plant | Carbuncles, | Toothache and worms in teeth and cough. |
| Bauhinia racemosa Lamk. | Leguminosae | Kanchan | Leaf | Dysurea | Leaves used in burning sensation of urine. |
| Boerhaavia diffusa Linn. | Nyctaginaceae | Sathi | whole plant | Eye, diarrheea | Used in urinary disorder, eyes. |
| Bombax ceiba Linn. | Bombacaceae | Semal | Flower, root | Burning sensation, leucorrhoea | Sexual diseases, burning sensation and sweta pradara |
| Borassus flabelifer Linn. | Palmae | Tad | Root | Contraceptive | To develop sterility and contraceptive |
| Boswellia serrata Colebr. | Burseraceae | Salar | Gum | Skin eruptions | Fragrant ointment and for impotency. |
| Bridelia retusa (Linn.) Spreng. | Euphorbiaceae | Lapano | Bark | Contraceptive | Bark is given orally to women to develop sterility and a |

| | | | | | |
|------------------------------|----------------|-----------------|-------------|------------------------------|---|
| | | | | | ctas contraceptive |
| Butea monosperma (Lam.) Taub | Leguminosae | Dhok, Khakhra | Whole plant | Leucorrhoea, cuts, dysentery | In leucorrhoea, excessive bleeding, sun stroke, pain delivery, Diarrhoea and hernia |
| Calotropis gigantea Linn. | Asclepiadaceae | Aak | Whole plant | Diarehoea, malaria | Latex used in Diarrhoea and burns. |
| Calotropis procera R. Br. | Asclepiadaceae | Akarda | Leaf | Abdominal pain | Tied on abdomen in abdominal pain |
| Capparis decidua Edgew | Capparidaceae | Kareera | Leaf, root | Eye disease | In eyes in corneal opacity, headache. |
| Capparis sepiaria Linn. | Capparidaceae | Himsra | Root | Cough | Powder of root is used in cough. |
| Cassia auriculata Linn. | Leguminosae | Alori | Leaf | Tuberculosis | Leaf extract is used in tuberculosis |
| Cassia fistula Linn | Leguminosae | Garmala | Seed, fruit | Menstrual disorder | Used in rakta pradar and diarrhoea. |
| Cassia tora Linn. | Leguminosae | Panvada | Leaf, seed | Dispepsia, | Used in night blindness & skin diseases. |
| Cayratia trifolia Domin. | Vitaceae | Khhata nimbi | Root | Abdominal Disorders | Applied on snake bite and carencules, diabetic, fractured and abdominal disorders. |
| Celosia argentea Linn. | Amaranthaceae | Kukidi | Seeds | Uterus diseases | Crushed seed is taken orally |
| Ceropegia bulbosa Roxb., | Asclepiadaceae | Khadula, Khadia | Seed, tuber | Stone, deafness, | To cure deafness, |

| | | | | | |
|--|----------------|-----------------|-----------------|---------------------------------------|--|
| Ceropegia Tuberosa Roxb. | | | | animal bite | urinary bladder stone, scorpion bite, fertility and vitality. |
| Chlorophytum borivillianum Baker | Liliaceae | Dholi musali | Tubers. | Enemic, fractured & leucorrhoea | To cure fractured bone, leucorrhoea and for male as tonic. |
| Chrozophora rottleri A. Juss. | Euphorbiaceae | Suballi | Leaves | Sun burn | Applied on the affected portion of body. |
| Cicer arietinum Linn. | Leguminosae | Chana | Seed | Rejuvenator, | Bread prepared of chana is taken in asthma. |
| Citrullus colocynthis Schrad. | Cucurbitaceae | Ingalboli | Fruit, seed | Abortifacient | Used for abortion and dental worm. |
| Citrus Medica Linn | Rutaceae | Bifora | Fruit | Dyspepsia | Fruit powder is used orally in dyspepsia. |
| Cleome viscosa Linn. | Capparidaceae | Kagiker | Whole plant | Fever | Juice of plant is dropped in the ear. |
| Clerodendrum phlomidis Linn. f. | Verbenaceae | Arani | Leaves | Joint pain | Applied on joints, in joint pain. |
| Clitoria ternatea Linn | Leguminosae | Vardia | Leaf | Eye disease | To cure swelling due to syphilis and eyes. |
| Cocculus sativus Linn. | Menispermaceae | Jalimani | Leaf | Leucorrhoea | To sterility in man and sweta pradar |
| Cocos nucifera Linn. | Palmae | Nariyal | Fruit, fibre | Leucorrhoea | Used as abortifacient and rakta pradar. |
| Colocasia esculenta (Linn.)Schott. | Araceae | Arbi | Tuber | Growth of hairs | Juice is used for good growth of |

| | | | | | |
|--|----------------|--------------|--------------|----------------------------|--|
| | | | | | hairs. |
| Commiphora wighti Linn. | Burseraceae | Gugal | Resin, latex | Urinary, wounds, Pyorrhea | Applied as an antiseptic, Infusion and on wounds |
| Convolvulus microphyllus Linn | Convolvulaceae | Bhumari | Whole plant | Constipation | Dried powder is taken orally with Water |
| Corallocarpus epigaeus (Rottl. & Willd) Hook | Cucurbitaceae | Mirchia Kand | Root | Diarrhoea, Stomachache | Used in stomach tumor or stomachache, typhoid and diarrhea. |
| Corchorus depressus (Linn.) Stocks | Tiliaceae | Baphuli | Whole plant | Menstrual, Polyurea | To gain sexual vitality, menstrual disorder and swap an dosh. |
| Cordia gharaf Roxb | Boraginaceae | Gundi | Bark, leaf | Toothache | Leaf is used in abortificiet. |
| Costus speciosus (Koen.) JESmith | Zingiberaceae | Maha lakdi | Root/ | Asthma, earache | To cure asthma, rheumatism and earache. |
| Crateva magna (Lour.) D.C. | Capparidaceae | Vaivana | Stem bark | Dysurea | To use orally in kidney stone and Abortion |
| Crinum asiaticum Linn. | Amaryllidaceae | Kanda' | Tuber, leaf | Ear ache, wound | Used for wound and in earache. |
| Curculigo orchioides Gaertn. | Amaryllidaceae | Kali musli | Root | Diarrhoea, syphilis | Used in gonorrhoea, syphilis, scorpion bite, leucorrhoea and Diarrhoea |
| Curcuma amada Roxb | Zingiberaceae | Amba haldi | Rhizome | Dropsy, snake bite, Asthma | To cure dropsy, asthma in cattle and snake bite, fractured bone. |
| Curcuma longa Linn. | Zingiberaceae | Haldi | Rhizome | Antifertility, | Used menstruation, |

| | | | | | |
|--------------------------------|----------------|--------------|-------------|-------------------------------|--|
| | | | | | Arthritis and elephantiasis. |
| Cuscuta hyalina Roth | Convolvulaceae | Amarbel | Whole plant | Abortifacient | Used orally for abortion and chest pain. |
| Cyamopsis tatarigoloba Taub. | Leguminosae | Ganwar | Leaf, seeds | Asthma swellings | Used in asthma and Swellings. |
| Datura innoxia Mill | Solanaceae | Dhatura | Leaf, seed | Asthma, malaria | Used in asthma, wound and malaria. |
| Daucus carota Linn | Umbelliferae | Gajar | Seed | Eye sight, sterility | To cure sterility in males, eyesight, to regularize menstruation and abortifacient |
| Dendrocalamus strictus (Roxb.) | Gramineae | Bans | Root | Dysurea | Paste of root is orally is used in dysurea |
| Desmostachya bipinnata Stapf. | Gramineae | Dabha | Whole plant | Fistula-in-ano | Ash of dhaba used in fistula-in-ano. |
| Dioscorea bulbifera Linn. | Dioscoreaceae | Jatashankari | Tuber | Contraceptive, abdominal pain | Used as contraceptive, abdominal pain, in asthma or cough and bone fracture. |
| Dolichandrone falcata Seem | Bignoniaceae | Milod | Leaf | Throat disease | Leaves are tried on affected part. |
| Dolichos biflorus Linn. | Leguminosae | Kulthi | Whole plant | Kidney stone | The plant is taken orally in kidney stone. |
| Annona squamosa Linn | Annonaceae | Sitaphal | Root | Dyspepsia | Used orally with mother's milk. |
| Eclipta Prostrata | Compositae | Bhanghra | Leaf | Wound | The leaf is locally applied |

| | | | | | |
|------------------------------------|---------------|--------|-------|-------------------|---|
| Linn. | | | | | over the skin. |
| Eulophia ochreatea Lindl | Orchidaceae | Mishri | Bulb | Diarrhoea | Crushed bulb used in diarrhoea. |
| Euphorbia caducifolia Haines | Euphorbiaceae | Thor | Latex | Body ache | Used as an abortifacient and body ache. |
| Euphorbia fusiformis Don. | Euphorbiaceae | Mooli | Tuber | Rheumatic pain | Paste of root applied in rheumatic pain. |

4. Source: Folk herbal medicines used by the Meena community in Rajasthan, Ajay Kumar Meena, M. M. Rao National Institute of Ayurvedic Pharmaceutical Research, CCRAS, Moti Bagh Road, Patiala 147001 (Punjab), India